



Fig. 2. Hypothetical mechanism for the synthesis of acid phosphatase f and s isozyme. N, F, S, C = ACP1 exons; i = ACP1 introns.

REFERENCES

- Dissing J (1986) Red cell acid phosphatase: only two different enzymes - the "slow" and the "fast" enzyme - determine different biochemical properties of the six common phenotypes. In: Brinkmann B and Henningsen K (eds) *Advances in Forensic Haemogenetics*, vol. 1. Springer, Berlin Heidelberg New York Tokyo, p 127
- Dissing J (1987) Immunochemical Characterization of Human Red Cell Acid Phosphatase Isozymes. *Biochem Genet* 25: 901-918
- Dissing J, Sensabaugh GF (1987) Human red cell acid phosphatase: evidence for differences in the primary structure of the two isozymes expressed by the ACP1*B allele. *Biochem Genet* 25: 919-927
- Dissing J, Svensmark O (Submitted for publication) Human red cell acid phosphatase: Purification and properties of the A, B and C isozymes.
- Fenton MR, Richardson KE (1971) Human erythrocyte acid phosphatase: resolution and characterization of the isozymes from three homozygous phenotypes. *Arch Biochem Biophys* 142: 13-21
- Harris H (1980) *The principles of Human Biochemical Genetics*, Elsevier/North-Holland, Amsterdam, New York Oxford, p 190
- Hopkinson, D.A., Spencer, N. and Harris, H. (1964). Genetical studies on human red cell acid phosphatase. *Am. J. Hum. Genet* 16: 141-154